

Comparison of the habitat conditions of peat-moss phytocoenoses dominated by *Eriophorum angustifolium* Honck. or *Carex rostrata* Stokes from mires in Western Poland

Maciej Gąbka^{1*}, Paweł M. Owsiany² & Tadeusz Sobczyński³

¹Department of Hydrobiology, Faculty of Biology, Adam Mickiewicz University, Umultowska 89, 61-614 Poznań, Poland, e-mail: gmaciej@amu.edu.pl

²Department of Geomorphology, Faculty of Geographical and Geological Science, Adam Mickiewicz University, Dziegielowa 27, 61-680 Poznań, Poland

³Department of Water and Soils Analysis, Faculty of Chemistry, Adam Mickiewicz University, Drzymały 24, 60-613 Poznań, Poland

Abstract: The presented phytosociological and habitat study concentrates on the moss phytocoenoses dominated by *Eriophorum angustifolium* or *Carex rostrata*, from 26 *Sphagnum* dominated peatlands adjacent to the lakes of the Wielkopolska region in Western Poland. Associations with these species dominate in the vegetation of the analysed mire configurations. 19 chemical parameters of surface water were analysed. Particular attention was paid to the determination of the habitat trophy and its indicators, the concentration of factors connected with the hardness of water, and the content of humic substances.

Phytocoenoses with *E. angustifolium* appeared in habitats where the analysed water parameters took lower values and more narrow amplitudes than those of the phytocoenoses with *C. rostrata*. The most important factors differentiating these two types of phytocoenoses, turned out to be the following: water colour, conductivity, Mg, Na and dissolved SiO₄. In conclusion, the appearance of phytocoenoses with *C. rostrata* was connected with habitats richer in nutrients and more advanced in the process of mineralisation than those of *E. angustifolium*.

Key words: *Carex rostrata*, *Eriophorum angustifolium*, mires vegetation, habitat conditions, habitat indicators, water chemistry, phytocoenoses

1. Introduction

Because of their transitional character and location between land and water, *Sphagnum* dominated peatlands play an essential role in the functioning of humic lakes. Physico-chemical character and biocenotic conditions of these lakes are conditioned by the constant flow of humic substances and nutrients provided by the direct catchment basin, and in particular by the zone of mires (e.g. Wojciechowski 1970, 1990; Górniak 1996, 1997; Górniak *et al.* 1999; Kruk 1997, 1999; Carpenter & Pace 1997; Williamson *et al.* 1999; Szymeja 2000; Wojtuń 2006).

Therefore, in order to determine the role of a given community in the accumulation, or in the transfer of particular elements of a habitat, it is essential to deter-

mine first the mutual relationships between habitat parameters of different phytocoenoses occupying the mires. The results of such a study may help to determine the role of vegetation in the diagnosis of abiotic conditions in mires and their impact on physico-chemical parameters of lake water (e.g. Wojciechowski 1990; Wojciechowski & Górniak 1990; Górniak 1996, 1997; Górniak *et al.* 1999; Kruk 1997, 1999; Gąbka *et al.* 2002, 2004).

The paper presents detailed characteristics of habitat conditions of phytocoenoses dominated by either *Eriophorum angustifolium* Honck. or *Carex rostrata* Stokes, located on peatlands adjacent to natural humic lakes. These species are considered characteristic of the associations (Brzeg *et al.* 2000; Brzeg & Wojterska

2001): *Sphagno recurvi-Eriophoretum angustifolii* Hueck 1925 *nom. invers. et nom. mut.* (syn. *Eriophoro angustifolii-Sphagnetum recurvi* M. Jasnowski *et al.* 1968 *em. Succow* 1974 *p.p. nom. illeg.*) and *Sphagno apiculati-Caricetum rostratae* Osvald 1923 *em. Steffen* 1931 (syn. *Sphagno-Caricetum rostratae* Steffen 1931 *em. Dierssen* 1982) from the *Scheuchzerio-Caricetea fuscae* (Nordhagen 1936) R.Tx. 1937 class. Despite their wide presence in the territory of Poland and Central Europe, these associations are relatively little known and ambiguously presented in literature (e.g. Succow 1974; Jasnowska & Jasnowski 1983a, 1983b; Borysiak *et al.* 1998; Brzeg *et al.* 1996; Timmermann 1993; Matuszkiewicz 2001). In the case of *Sphagnum* dominated peatlands ecosystems of western Poland, these associations are the basic element of the mire vegetation. The aim of the study was to determine the differences between the ecological amplitudes of phytocoenoses dominated by *E. angustifolium* and those with *C. rostrata*.

2. Materials and methods

2.1. Study area

For the analysis, 26 *Sphagnum* dominated peatlands were selected (Table 1), all of them adjacent to the humic lakes distributed in the lake districts of the Wielkopolska region and of the Southern Pomerania (Kondracki 1998), both in western Poland. These mires are placed around small natural lakes (<10 ha), mostly very shallow

(<5 m) and in the majority of cases situated in forests with the domination of a pine forest in a drainage basin. As far as the appearance of this type of ecosystems in Poland is concerned, these lakes are a bit off-centre, being mostly associated with the Pomerania region (e.g. Hutorowicz *et al.* 1999; Kraska *et al.* 1999). In the analysed region, such bog-lake ecosystems are a very rare element of the lakeland landscape.

The analysed region is characterised by small annual precipitation, ca. 500 mm per year. Mean annual temperatures range from 7.2°C in the north-eastern part of the region to 8.2°C in the west. The coldest season coincides with January (-1.5 – -2.5°C), the warmest weather is, over the years, traditionally noted in July (17.5 – 19.0°C) (Woś 1994).

2.2. Vegetation data

Phytosociological and habitat observation concerned mire phytocoenoses dominated by *Eriophorum angustifolium* or *Carex rostrata*. Phytosociological specificity of patches classified as *Sphagno recurvi-Eriophoretum angustifolii* and *Sphagno apiculati-Caricetum rostratae* was based on considerable density and abundance of characteristic species, i.e., *E. angustifolium* and *C. rostrata*, respectively.

Field investigations were carried out in the summer during the period of maximal growth (July, August) between 2001-2004. Phytosociological relevés were taken according to the Braun-Blanquet method (Braun-Blanquet 1951). Altogether, 18 patches with *Eriophorum*

Table 1. The location of mires with patches of *Eriophorum angustifolium* (EA) and *Carex rostrata* (CR) and number of relevés investigated in the study

No.	Lakes	Administrative unit	Geographical coordinates	EA	CR
1	Święte	Wieleń	N 52°48'5'' E 16°12'3''	.	1
2	Moczydło	Drawsko	N 52°49'8'' E 16°00'0''	.	2
3	Perskie	Drawsko	N 52°45'6'' E 16°01'8''	.	1
4	Rzecińskie	Wronki	N 52°45'7'' E 16°45'7''	1	1
5	Pokraczyn	Wronki	N 52°46'2'' E 16°21'2''	2	.
6	Pustelnik I	Wronki	N 52°46'7'' E 16°19'1''	3	.
7	Pustelnik II	Wronki	N 52°46'8'' E 16°19'1''	3	.
8	Wilcze Błoto	Wronki	N 52°49'4'' E 16°22'8''	1	2
9	Mnich Mały	Sieraków	N 52°39'0'' E 16°01'9''	1	.
10	Zamorze	Pniewy	N 52°30'6'' E 16°12'6''	.	1
11	Skrzynka	Mosina	N 52°15'3'' E 16°47'2''	.	2
12	Niewiemko	Budzyń	N 52°53'7'' E 16°45'8''	.	1
13	Czarne Małe	Murowana Goślina	N 52°31'8'' E 17°05'3''	.	1
14	Kańskie	Powidz	N 52°24'8'' E 18°01'5''	.	1
15	Czarne	Witkowo	N 52°28'6'' E 17°53'4''	.	1
16	Święte	Przemęt	N 51°57'7'' E 16°46'3''	2	.
17	Żurawin	Rychwał	N 52°02'0'' E 18°05'5''	1	.
18	Modre	Sępólno Krajeńskie	N 53°25'6'' E 17°20'3''	1	.
19	Czarne	Kaczory	N 53°08'1'' E 16°55'2''	.	1
20	Smolary	Szydłowo	N 53°17'8'' E 16°44'0''	1	.
21	Kuźnik Mały	Piła	N 53°11'9'' E 16°44'4''	.	1
22	Kuźniczek	Piła	N 53°11'5'' E 16°44'3''	.	2
23	Kuźnik Bagienny	Szydłowo	N 53°12'5'' E 16°44'0''	1	.
24	Kuźnik Olszowy	Szydłowo	N 53°12'4'' E 16°43'5''	1	1
25	Czapla	Dobrzyca	N 53°11'3'' E 16°48'1''	.	1
26	Okoniowe	Piła	N 53°11'1'' E 16°48'2''	.	2

angustifolium from 12 mires and 22 patches with *Carex rostrata* from 17 mires were sampled (Table 1). The floristic composition under study was collected in one synthetic table of shortened form. Abundance, constancy and coefficient of cover were worked out. Abundance was determined according to a modified six-degree scale by Braun-Blanquet. Constancy was determined according to the following scale on the basis of species abundance: V – species occurring in 80.1-100% of relevés taken in the phytocoenoses; IV – species present in 60.1-80.0%; III – species present in 40.1-60.0%; II – species present in 20.1-40.0%; I – species present in 10.1-20.0%; + – species present in 5.1-10.0% and $r \leq 5.0\%$.

Vascular plants names are those used by Mirek *et al.* (2002), of mosses by Ochyra *et al.* (2003), and the nomenclature of the syntaxa follows Brzeg and Wojterska (2001).

2.3. Environmental data

For each of the phytocoenoses in which phytosociological relevé was made, water samples were taken, too, from the central parts of each patch (Kłosowski & Tomaszewicz 1993; Kłosowski 1999). Each of the analysed water samples consisted of three sub-samples taken from the patches up to 0.5 m deep. The samples were taken from shallow piezometric wells in the centre of each patch. Water samples were poured into three plastic containers, two of which were immediately preserved: one by addition of 1 ml of nitrate acid, the other with 1 ml of 95% chloroform; the third sample was not treated with any preservatives. Water pH and temperature, conductivity, dissolved oxygen and water oxygenation were measured directly on the site; the remaining parameters were measured after the samples had been taken to the laboratory.

Physical and chemical analysis were performed according to the Siepak (1992) and Hermanowicz *et al.* (1998) methods. 19 physical-chemical parameters of water were analyzed: colors (platinum-cobalt standard method), pH (with pH-meter), dissolved organic carbon (DOC, with organic carbon analyzer), conductivity (electrolytic method), NH_4^+ (by Nessler's colorimetric method), O_2 dissolved, saturation with O_2 (electrometric method), NO_3^- (by cadmium reduction method), PO_4^{3-} (by the colorimetric ascorbic acid method), SO_4^{2-} (by nephelometric method), Cl⁻ (by Mohr's argentometric method), total hardness, carbonate hardness, Ca^{2+} , Mg^{2+} (wersenian method), SiO_2 dissolved (heteropoly blue method), Na^+ , K^+ and total Fe (by atomic absorption spectrometry method).

In order to compare the habitats, average values and ecological amplitudes of particular water parameters were taken into consideration. The significance of differences in water properties between groups of samples from the sites dominated by *Eriophorum angustifolium* and by *Carex rostrata* was tested by Mann-Whitney *U* test.

3. Results

3.1. Phytosociological characteristics of the phytocoenoses

Phytosociological analysis of phytocoenoses dominated by *Eriophorum angustifolium* and by *Carex rostrata* suggests that these species often tend to form dense compact stands (cover-abundance coefficient mostly 4). The species composition and phytosociological data are contained in Table 2. Phytocoenoses whose herb layer was dominated by *C. rostrata* were floristically richer than those with *E. angustifolium*. Patches of the analysed associations were clearly marked with the constant presence of species belonging to the class *Oxycocco-Sphagnetea*. In dryer conditions, the share of shrubs and low trees was visibly greater especially *Pinus sylvestris* and *Betula pubescens*.

In the observed group of phytocoenoses with *E. angustifolium* (Table 2, EA) the most constant taxa appear to be such species as *Carex limosa* and *Menyanthes trifoliata*. Among vascular plants which were present in the patches, the most numerous was *Rhynchospora alba* (III constancy class). The moss layer was dominated mostly by *Sphagnum fallax* (V constancy class) and less often by *Sphagnum magellanicum* (III constancy class). In several patches only it was *Sphagnum flexuosum* that exhibited the greatest coverage in the moss layer. In a number of analysed phytocoenoses with *E. angustifolium* we could notice a considerable participation of other mosses of the classes *Oxycocco-Sphagnetea* and *Scheuchzerio-Caricetea fuscae*, i.e. *Aulacomnium palustre*, *Polytrichum strictum*, *Calliergon stramineum* and *Warnstorfia exannulata*.

In the patches dominated by *Carex rostrata* most constant were species of the class *Scheuchzerio-Caricetea fuscae*, such as *Menyanthes trifoliata*, *Comarum palustre* and *Eriophorum angustifolium*, then the class *Oxycocco-Sphagnetea*. Relatively larger, too, was the share of *Thelypteris palustris* and *Lysimachia vulgaris* (III constancy class), as well as some species with constancy II (Table 2, CR). The moss layer, in turn, was most often dominated by *Sphagnum fallax* (V constancy class); noticeable, too, was a higher constancy of such species as *Sphagnum palustre* and *Calliergon stramineum*. In the structure of these phytocoenoses, larger participation of tree species was visible, mainly of *Betula pubescens*, than it was in the case of phytocoenoses with *Eriophorum angustifolium*.

Phytocoenoses dominated by *Eriophorum angustifolium* or *Carex rostrata* were the most important peat-forming communities in the analysed mires. In the beltlike configuration of vegetation, phytocoenoses with *E. angustifolium* were either adjacent to the lakes water table, or they were preceded by patches of *Caricetea limosae* (Beger 1922) Osvald 1923 em. Dierßen 1982,

Table 2. A synoptic table of the vegetation data studied. EA – community of *Eriophorum angustifolium*, CR – community of *Carex rostrata*

Community type	EA	CR
Number of relevés	18	22
Number of mires	13	22
Average relevé area [m ²]	29	28
Total number of species	59	88
Mean number of species in one relevé	15	18
I. Dominant species		
<i>Eriophorum angustifolium</i>	V ³⁻⁴	IV ⁺¹
<i>Carex rostrata</i>	IV ^{r-2}	V ³⁻⁴
II. Characteristic species of the order <i>Scheuchzerietalia palustris</i>		
<i>Sphagnum fallax</i>	V ³⁻⁵	V ¹⁻⁵
<i>Carex limosa</i>	IV ⁺¹	II ⁺
<i>Rhynchospora alba</i>	III ⁺¹	r ⁺
<i>Scheuchzeria palustris</i>	I ⁺¹	r ⁺
<i>Carex lasiocarpa</i>	I ⁺¹	I ⁺¹
<i>Sphagnum flexuosum</i>	I ¹⁻³	II ⁺⁴
<i>Sphagnum teres</i>	r ²	II ¹⁻²
<i>Carex diandra</i>	r ⁺	I ⁺¹
<i>Sphagnum warnstorffii</i>	r ⁺	I ⁺¹
III. Characteristic species of the order <i>Caricetalia fuscae</i>		
<i>Carex canescens</i>	II ⁺	.
<i>Carex echinata</i>	II ⁺¹	I ⁺
<i>Agrostis canina</i>	II ⁺¹	II ⁺¹
<i>Carex nigra</i>	I ⁺	.
<i>Epilobium palustre</i>	I ⁺	II ^{r+}
<i>Calamagrostis stricta</i>	+ ⁺¹	II ^{r-1}
<i>Hydrocotyle vulgaris</i>	r ⁺	II ⁺
IV. Characteristic species of the class <i>Scheuchzerio-Caricetea fuscae</i>		
<i>Menyanthes trifoliata</i>	IV ^{r-1}	IV ^{r-3}
<i>Calligon stramineum</i>	IV ⁺²	IV ⁺²
<i>Warnstorfia exannulata</i>	II ⁺¹	.
<i>Comarum palustre</i>	II ^{r-1}	III ^{r+}
<i>Calla palustris</i>	r ⁺	+ ^{r+}
<i>Sphagnum riparium</i>	.	+ ³⁻⁴
V. Characteristic species of the class <i>Oxycocco-Sphagnetea</i>		
<i>Drosera rotundifolia</i>	V ⁺	IV ⁺¹
<i>Oxycoccus palustris</i>	V ⁺⁴	III ⁺³
<i>Aulacomnium palustre</i>	IV ⁺¹	III ⁺²
<i>Sphagnum magellanicum</i>	III ¹⁻³	I ⁺¹
<i>Eriophorum vaginatum</i>	II ⁺²	II ⁺
<i>Polytrichum strictum</i>	II ⁺¹	III ⁺²
<i>Sphagnum capillifolium</i>	r ⁺	I ⁺²
<i>Andromeda polifolia</i>	r ⁺	I ⁺¹
VI. Others		
<i>Pinus sylvestris</i> juv.	IV ⁺	II ^{r+}
<i>Pinus sylvestris</i> fo. <i>turfosa</i> b	III ⁺²	III ⁺²
<i>Juncus effusus</i>	II ^{r+}	II ⁺²
<i>Sphagnum denticulatum</i>	II ⁺¹	r ¹
<i>Betula pubescens</i> b	II ⁺²	IV ⁺³
<i>Betula pubescens</i> juv.	II ^{r+}	III ^{r+}
<i>Sphagnum palustre</i>	II ⁺²	III ⁺²
<i>Lysimachia thyrsoflora</i>	II ^{r+}	I ⁺
<i>Lycopus europaeus</i>	I ^{r+}	II ^{r+}
<i>Molinia caerulea</i>	I ^{r+}	+ ⁺
<i>Calliergonella cuspidata</i>	I ⁺	I ⁺¹
<i>Lysimachia vulgaris</i>	I ⁺	III ⁺
<i>Thelypteris palustris</i>	r ^r	III ⁺¹
<i>Salix aurita</i> b	r ⁺	II ⁺¹
<i>Frangula alnus</i> b	r ⁺	I ⁺¹
<i>Frangula alnus</i> juv.	r ^r	I ⁺
<i>Typha latifolia</i>	r ¹	I ^{r+}
<i>Salix cinerea</i> b	r ⁺	+ ⁺
<i>Carex disticha</i>	r ⁺	r ⁺
<i>Phragmites australis</i>	.	II ⁺
<i>Peucedanum palustre</i>	.	II ^{r+}
<i>Alnus glutinosa</i> b	.	II ⁺¹
<i>Alnus glutinosa</i> juv.	.	II ^{r+}
<i>Dryopteris cristata</i>	.	II ^{r+}
<i>Galium palustre</i>	.	II ^{r+}

Sphagno tenelli-Rhynchosporium albae Osvald 1923 nom. invers. or *Caricetum lasiocarpae* Osvald 1923. In turn, phytocoenoses with *Carex rostrata* most often came either next to the patches with *Thelypterido-Phragmitetum* Kuiper 1958 or *Cladietum marisci* Allorge 1922 ex Zobrist 1935 on the water side, or developed behind other *Sphagnum* communities.

3.2. Habitat conditions of the phytocoenoses

Phytocoenoses with *Eriophorum angustifolium* were connected with complex peat habitats, usually highly hydrated, yet, without the stagnant water in the patches. In contrast, most of the patches with *Carex rostrata* showed low levels of ground water.

Differences in the physico-chemical parameters of water between the two types of analysed phytocoenoses were observed in the case of most of the parameters, especially those connected with the hardness of water, and with the concentration of humic substances and nutrients. One of the characteristic features of the habitats of the analysed species is the high content of nutrients and considerable habitat oxygenation, higher in the patches dominated by *Carex rostrata*. Yet, no significant difference between the habitats of the two communities was noted as far as the water pH is concerned. In the case of the habitats with *Carex rostrata*, the pH value had a wider range and higher maximum than in the case of the habitats with *Eriophorum angustifolium*.

Standardised data on all the water parameters suggest that most of the phytocoenoses with *E. angustifolium* exhibited low range of values (Fig. 1). Only the average values of specific acidity (pH) and PO_4^{3-} exceeded the line of general average values. Groups of phytocoenoses dominated by the analysed species developed in wide ranges of water characteristics; their values, however, were considerably lower in the case of phytocoenoses with the domination of *E. angustifolium*.

The water in the phytocoenoses with *E. angustifolium* exhibited lower values of the colour of water and conductivity; moreover, in comparison to the patches with *C. rostrata*, it was also poorer in Ca^{2+} , Na^+ , K^+ , NO_3^- , N-NH_4^+ and dissolved SiO_2 . Yet, the habitats with *E. angustifolium* were more abundant as far as the content of PO_4^{3-} and total Fe is concerned. With reference to such parameters as water colour, specific acidity (pH),




Fig. 1. The water properties in the habitat of phytocoenoses dominated by *Eriophorum angustifolium* (A) and *Carex rostrata* (B). Median value, lower and upper quartile based on the standardized data

Explanations: 1 – colors, 2 – pH, 3 – saturation with O_2 , 4 – O_2 dissolved, 5 – conductivity, 6 – N-NH_4^+ , 7 – N-NO_3^- , 8 – P-PO_4^{3-} , 9 – total Fe, 10 – total hardness, 11 – carbonate hardness, 12 – Ca^{2+} , 13 – Mg^{2+} , 14 – Na^+ , 15 – K^+ , 16 – SO_4^{2-} , 17 – Cl^- , 18 – SiO_2 dissolved, 19 – DOC

general hardness, Mg^{2+} and dissolved SiO_2 , in the case of the phytocoenoses with *C. rostrata* they were shifted above the general average values (Fig. 1). Waters of these patches were richer in the compounds of Ca^{2+} and Mg^{2+} , as well as in SO_4^{2-} and Cl^- . Especially wide amplitudes of values were visible in the case of parameters connected with the concentration of humic substances (colour, DOC), water fertility (NH_4^+) and habitat mineralization (conductivity, saturation with O_2 and O_2 dissolved).

Most important differences between water parameters of phytocoenoses with *Eriophorum angustifolium* and those of *Carex rostrata* were observed as far as the colour of water and conductivity are concerned, as well as the concentration of Mg^{2+} , K^+ and dissolved SiO_2 (Table 3). These parameters essentially differentiated the phytocoenoses with *E. angustifolium* from those with *C. rostrata*.

Sporadic species: II. *Sphagnum subnitens* SCR (r^+); *Sphagnum obtusum* SCR (r^2); III. *Triglochin palustre* SCR (r^+); *Ranunculus flammula* SCR (r^+); *Veronica scutellata* SCR (I^{++}); *Viola palustris* SCR (I^+); *Eleocharis quinqueflora* EA (r^+), SCR (r^+); *Sphagnum fimbriatum* SCR ($+^2$); V. *Polytrichum commune* SCR (r^+); *Sphagnum fuscum* EA (r^+); *Sphagnum angustifolium* EA (r^1); VI. *Picea exelsa* b SCR ($+^+$); *Pinus sylvestris* fo. *turfosa* a SCR (I^2); *Vaccinium myrtillus* SCR (r^+); *Plagiomnium elatum* EA (r^+); *Salix aurita* juv. SCR (II^{+1}); *Alnus glutinosa* a SCR ($+^+$); *Empetrum nigrum* SCR (r^1); *Equisetum fluviatile* SCR ($+^{+1}$); *Potentilla erecta* SCR ($+^+$); *Stellaria palustris* SCR (r^1); *Quercus robur* juv. SCR (I^{++}); *Juncus inflexus* SCR (r^+); *Salix repens* subsp. *rosmarinifolia* juv. SCR (r^2); *Betula pubescens* a SCR (I^{1-2}); *Ledum palustre* SCR ($+^{+1}$); *Dryopteris cartusiana* SCR (r^+); *Cladium mariscus* SCR (r^+); *Typha angustifolia* EA (r^+); *Polytrichum longisetum* EA (r^1), SCR (r^1); *Bidens cernua* EA (r^+); *Utricularia minor* EA (r^1); *Lythrum salicaria* SCR ($+^+$); *Scutellaria galericulata* EA (r^+); *Epilobium parviflorum* SCR ($+^{++}$); *Salix repens* subsp. *repens* b SCR (r^+); *Sphagnum squarosum* SCR (r^1); *Salix repens* subsp. *rosmarinifolia* SCR ($+^2$); *Cirsium palustre* (r^+); *Utricularia intermedia* (r^+)

Table 3. Physical and chemical properties of water found in habitats of phytocoenoses studied. Ranges and mean values and the significance of differences in water properties between groups phytocoenoses obtained from Mann-Whitney *U* test

Property		EA		SCR		Significance EA-SCR
		Range	Mean	Range	Mean	
Colour	mg Pt/l	37-469	191.44	79-535	288.86	+
DOC	mg C/l	24.49-121.47	54.2	15.04-102.39	51.16	ns
pH		3.4-4.9	4.36	3.52-5.94	4.45	ns
O ₂ dissolved	mg O ₂ /l	0.51-10.1	2.44	0.64-20	3.65	ns
Saturation with O ₂	%	6.9-126	28.60	6.43-200	39.71	ns
Conductivity	µS/cm	33.3-413	104.35	40.3-369	134.88	+
N-NH ₄ ⁺	mg N/l	0.47-3.2	2.12	0.4-5.1	2.44	ns
N-NO ₃ ⁻	mg N/l	0.1-1.5	0.55	0.1-2.1	0.67	ns
P-PO ₄ ³⁻	mg PO ₄ /l	0.24-5.16	1.94	0-5.12	1.33	ns
Total Fe	mg Fe/L	0.01-1.07	0.33	0.01-1.45	0.4	ns
Total hardness	mval/l	0.02-0.6	0.26	0.03-0.85	0.33	ns
Carbonate hardness	mval/l	0.01-0.5	0.14	0.01-0.9	0.24	ns
Ca ²⁺	mg Ca/l	0.26-7.43	2.36	0.26-13.21	4.1	ns
Mg ²⁺	mg Mg/l	0.09-3.17	1.156	0.26-4.03	1.861	+
Na ⁺	mg Na/l	0.27-6.29	1.95	0.4-23.4	4.43	+
K ⁺	mg K/l	1.46-19.12	7.58	0.47-21.4	8.05	ns
SO ₄ ²⁻	mg SO ₄ /l	0-34	1.94	0-35	4.64	ns
SiO ₂ dissolved	mg SiO ₂ /l	0.01-1.72	0.55	0.17-2.90	1.11	+
Cl ⁻	mg Cl/l	3-16	6.4	4-62	10.59	ns

Explanations: EA – community of *Eriophorum angustifolium* (n=18), CR – community of *Carex rostrata* (n=22), ns – non-significant

4. Discussion

Analyses of habitat requirements of the species, and their communities, occupying mires suggest mutual relationships between the species and the parameters of their habitats (e.g. Hájková & Hájek 2004a, 2004b; Navrátilová & Navrátil 2005; Hájková *et al.* 2002). Ground water level, reaction and conductivity have been regarded as the most important abiotic factors influencing the appearance of species and their communities in mires (e.g. Miserere *et al.* 2003; Tahvanainen & Tuomala 2003; Hájková & Hájek 2004a, 2004b). Numerous studies have suggested that species diversity and the presence of specific associations could be used as indicators of environmental characteristics and of anthropogenic impact (e.g. Hájková & Hájek 2004a, 2004b; Bragazza *et al.* 2005; Navrátilová & Navrátil 2005). Sometimes it is noticed, too, that regional habitat differences, observable in the case of the vegetation of waterlogged areas, may to some extent limit the indicative value of particular associations (e.g. Pelechaty 2004).

The analysed phytocoenoses dominated by *Eriophorum angustifolium* or *Carex rostrata*, which form the most important parts of the vegetation of the observed mires, were connected with different habitat requirements. It might suggest their bioindicative value. Phytocoenoses with *E. angustifolium* appear in habitats characterised by lower values of most of the analysed water parameters than those of phytocoenoses with *C. rostrata*. The most important elements differentiating these two groups of analysed phytocoenoses were the following water parameters: colour, conductivity, Mg²⁺, Na⁺ and dissolved SiO₂. In comparison to phytocoenoses

with *C. rostrata*, those with *E. angustifolium* were connected with the waters poorer in Ca²⁺, with smaller concentration of NH₄⁺ and NO₃⁻, and higher content of PO₄³⁻. The analysed habitats with *C. rostrata* were more fertile both as far as the content of nutrients is concerned, and with regard to such substances as Na⁺, K⁺, Ca²⁺ and SO₄²⁻.

It has to be pointed out that in the case of the analysed phytocoenoses of peat-bogs adjacent to the humic lakes in the Wielkopolska region, the water reaction was not an essential differentiating factor. The features analysed above can be regarded as characteristic for communities dominated by these mires species and present in the bog-lake ecosystems of western Poland.

The analysed habitats were characterised by wide ranges of ecological optima. They appeared in wide ranges in the case of acidic reaction of water acidity? (especially patches with *C. rostrata*), considerably coloured and rich in nutrients. Yet, some data present in the literature on the subject suggest the predilection of phytocoenoses with domination *E. angustifolium* or *C. rostrata* for habitats with moderate or slightly higher nutrient abundance (Matuła *et al.* 1997). In the peat-bogs of the Sudety Mountains, in the patches with these species a high concentration of nutrients was found, especially those of NO₃⁻ and NH₄⁺ (Matuła *et al.* 1997). We have to bear in mind, however, that patches of the analysed phytocoenoses of the bog-lake ecosystems had their *Sphagnum* substratum less hydrated than it is described by the data found in literature (Pietsch 1976; Brzeg *et al.* 1996; Matuła *et al.* 1997). Smaller water volume in the habitats, together with the processes of *Sphagnum* peat mineralisation in the peak of the vegetative season, are reflected in the characteristics of

water. In numerous patches, especially in those dominated by *C. rostrata*, high values were measured for such water parameters as: conductivity, oxygen concentration and oxygenation, or NH_4^+ ; together with low reaction, they all indicate the state of advanced mineralisation.

Wide ecological scale and the multi-facial character of communities created by *Carex rostrata* and by *Eriophorum angustifolium* have been pointed out in numerous studies (e.g. Hadač & Váňa 1967; Jasnowska & Jasnowski 1983a, 1983b; Matuła *et al.* 1997; Borysiak *et al.* 1998).

Most of the parameters, both in the case of phytocoenoses with *E. angustifolium* and those with *C. rostrata*, exhibited wide ranges of the occurrence. This fact may be connected with the domination of *Sphagnum fallax* in the moss layer of majority of the patches. We have to point out, though, that the habitat requirements and the diversity of ecological gradients in the case of *Sphagnum* peat mosses are different from those of vascular plants growing over peat-bogs (Hájková & Hájek 2004a, 2004b).

Sphagnum fallax is sometimes regarded as an indicator of advanced habitat trophism and particularly high concentration of N (Limpens *et al.* 2003). We may point to the important habitat-creating role of this dominating peatmoss, characteristic for wide expansion at the present state of habitat development (Jasnowska & Jasnowski 1983a, b; Limpens *et al.* 2003; Miserere *et al.* 2003). It has to be pointed, too, that the peat-bogs exhibiting a dense structure of *Sphagnum* peat, with *Sphagnum fallax* dominating in the moss layer, were very frequent in the analysed lake-bog ecosystems. Small floristic diversity of moss layer in the case of peat-bogs with *Sphagnum fallax*, despite the diversity of vascular species forming physiognomically separate patches within the limits of some *Sphagnum* bog objects, has

been described by Jasnowska *et al.* (2000) and Hájková and Hájek (2004a, 2004b).

In the mosses layer of phytocoenoses dominated by *E. angustifolium* we could locally distinguish a greater participation of peatmoss *Sphagnum flexuosum*. This taxon is considered in Poland as a species regionally characteristic for the association *Sphagno recurvi-Eriophoretum angustifolii* (Brzeg *et al.* 1996; Borysiak *et al.* 1998), and as an indicator of waterlogged habitats (Borysiak *et al.* 1998). However, phytocoenoses with *E. angustifolium* and a considerable participation of *Sphagnum flexuosum* were rarely met in the mires of humic lakes. Indirectly, this can suggest the overdrying of organogenic habitats.

On numerous occasions, in the patches with analysed species, especially with *Carex rostrata*, we could meet low trees and shrubs. This fact may be connected with natural dynamic transformations of mires into forest communities, conditioned by small precipitation in the analysed region.

The diagnosis of habitat conditions of mires dominated by phytocoenoses with *Eriophorum angustifolium* or *Carex rostrata* in the lake-bog ecosystems may be of help in the determination of physico-chemical condition of lake waters. The appearance of patches dominated by *C. rostrata* was observed mainly in the case of lakes with humotrophic waters, fertile and rich in nutrients. Phytocoenoses with *C. rostrata*, in turn, tended to appear optimally in mires adjacent to lakes of strongly marked dystrophic character (Gąbka *et al.* 2004, 2005).

Acknowledgements. Scientific work financed from the resources earmarked for science in years 2001-2003 as the Research Project no. 6 P04F 037 29. We thank to Dr. Barbara Fojcik (University of Silesia) for help in identification of mosses. The anonymous reviewers are thanked for their helpful comments.

References

- BORYSIK J., MELOSIK I. & STACHNOWICZ W. 1998. Szata roślinna i ochrona torfowiska przejściowego "Gogulec" koło Poznania. *Bad. Fizjogr. Pol. Zach., seria B-Botanika* 47: 159-175.
- BRAGAZZA L., RYDIN H. & GERDOL R. 2005. Multiple gradients in mire vegetation: a comparison of Swedish and an Italian bog. *Plant Ecology* 177: 223-236.
- BRAUN-BLANQUET J. 1951. *Pflanzensoziologie*. 631 pp. Springer Verlag, Wien.
- BRZEG A., KUŚWIK H., MELOSIK I. & URBAŃSKI P. 1996. Flora i roślinność projektowanego rezerwatu przyrody "Zielone Bagna" w Drawieńskim Parku Krajobrazowym. *Bad. Fizjogr. Pol. Zach. seria B-Botanika* 45: 121-145.
- BRZEG A., KUŚWIK H. & WYRZYKIEWICZ-RASZEWSKA M. 2000. Szata roślinna projektowanego rezerwatu "Ciświckie Bagna" koło Grodzca we wschodniej Wielkopolsce. *Roczniki AR w Poznaniu* 322: 21-67.
- BRZEG A. & WOJTERSKA M. 2001. Zespoły roślinne Wielkopolski, ich stan poznania i zagrożenie. In: M. WOJTERSKA (ed.). *Szata roślinna Wielkopolski i Pojezierza Południowopomorskiego. Przewodnik sesji terenowych* 52. Zjazdu PTB, pp. 39-110. Bogucki Wyd. Nauk., Poznań.
- CARPENTER S. R. & PACE M. L. 1997. Dystrophy and eutrophy in lake ecosystems: implications of fluctuating input. *Oikos* 78: 3-14.
- GĄBKA M., OWSIANNY P. M. & SOBECZYŃSKI T. 2004. Acidic lakes in the Wielkopolska region – physico-chemical properties of water, bottom sediments and the aquatic micro- and macrovegetation. *Limnological Review* 4: 81-88.

- GĄBKA M., OWSIANNY P. M., SOBCZYŃSKI T. & ZIOŁA A. 2005. The spring-lake Niewiemko – horizontal diversity of algae and macrophytes communities on the background of habitat conditions (Poland). *Limnological Review* 5: 75-80.
- GĄBKA M., PEŁECHATY M. & MATUSZAK K. 2002. Phytocoenotic and floristic differentiation among the chosen lake-bog ecosystems of the Wielkopolska region. *Acta Agrophysica* 67: 77-83.
- GÓRNIAK A. 1996. Substancje humusowe i ich rola w funkcjonowaniu ekosystemów słodkowodnych. 155 pp. Wyd. Filii UW WB, Białystok.
- GÓRNIAK A. 1997. Kwasy humusowe a eutrofizacja ekosystemów wodnych. In: *Zastosowanie biotechnologii ekosystemalnych do poprawy jakości wód*. Zeszyty Naukowe Komitetu "Człowiek i Środowisko" 18: 101-114.
- GÓRNIAK A., JEKATIERYN CZUK-RUDCZYK E. & DOBRZOŃ P. 1999. Hydrochemistry of three dystrophic lakes in north-eastern Poland. *Acta hydrochim. hydrobiol.* 27(1): 12-18.
- HADAČ E. & VÁŇA J. 1967. Plant Communities of Mires in the Western Part of the Krkonoše Mountains, Czechoslovakia. *Folia Geobot. Phytotax.* 2(2): 213-254.
- HÁJKOVÁ P. & HÁJEK M. 2004a. Bryophyte and vascular plant response to base-richness and water level gradients in Western Carpathian *Sphagnum*-rich mires. *Folia Geobotanica* 39: 335-351.
- HÁJKOVÁ P. & HÁJEK M. 2004b. *Sphagnum*-mediated successional pattern in the mixed mire in the Muránska plainina Mts (Western Carpathians, Slovakia). *Biologia, Bratislava* 59(1): 65-74.
- HÁJKOVÁ P., HEKERA P. & HÁJEK M. 2002. Spring fen vegetation and water chemistry in the western Carpathian flysch zone. *Folia Geobotanica* 37: 205-224.
- HERMANOWICZ W., DOZAŃSKA W., DOJLIDO J. & KOZIOROWSKI B. 1999. Fizyczno-chemiczne badania wody i ścieków. 847 pp. Arkady, Warszawa.
- HUTOROWICZ A., TUNOWSKI J. & ZDANOWSKI B. 1999. Dystroficzne jeziora Wigierskiego Parku Narodowego – struktura, funkcjonowanie i zagrożenia. In: S. RADWAN & R. KORNIJOW (eds.). *Problemy aktywnej ochrony ekosystemów wodnych i torfowiskowych w polskich parkach narodowych*, pp. 219-230. Wyd. UMCS, Lublin.
- JASNOWSKA J. & JASNOWSKI M. 1983a. Zbiorowiska roślinne związku *Caricion lasiocarpae* V. d. Bergh. ap. Lebr. 1949 torfowisk mszarnych na Pojezierzu Bytowskim. *Zesz. nauk. AR w Szczecinie, Rolnictwo* 32, Seria Przyrodnicza 104: 65-74.
- JASNOWSKA J. & JASNOWSKI M. 1983b. Roślinność rzędu *Caricetalia fuscae* (= *nigrae*) Nordh. 36 emend. Preis. ap. Oberd. 49 torfowisk mszarnych Pojezierza Bytowskiego. *Zesz. nauk. AR w Szczecinie, Rolnictwo* 32, Seria Przyrodnicza 104: 81-87.
- JASNOWSKA J., KOWALSKI W. & MARKOWSKI S. 2000. Rezerwat przyrody "Mszar nad jeziorem Piaski" na Wysoczyźnie Łobeskiej. *Folia Univ. Agric. Stetin.* 213, *Agricultura* 85: 105-128.
- KŁOSOWSKI S. 1999. Synecological studies on littoral vegetation in northern Poland. *Acta Hydrobiol.* 41 Suppl. 6: 49-54.
- KŁOSOWSKI S. & TOMASZEWICZ H. 1993. Standortverhältnisse der Gesellschaften mit Dominanz einzelner Nymphaeaceen in Nordeost-Polen. *Tuexenia* 13: 75-90.
- KONDRACKI J. 1998. *Geografia regionalna Polski*. 440 pp. Wyd. Nauk. PWN, Warszawa.
- KRASKA M., PIOTROWICZ R. & RADZISZEWSKA R. 1999. Dystrophication as the chief factor of changes in the physico-chemical properties of water and vegetation of lobelian lakes of the Bory Tucholskie National Park (NW Poland). *Acta Hydrobiol.* 41, Suppl. 6: 127-135.
- KRUK M. 1997. Element retention and loss by hydrologically modified peatlands in the Mazurian Lakeland (North-Eastern Poland) II. Peatland with eutrophicated transition bog. *Pol. Ecol. Stud.* 23(1-2): 23-44.
- KRUK M. 1999. Peatlands – barrier or source of phosphorus inflow to lakes? *Acta Hydrobiol.* 41, Suppl. 6: 73-81.
- LIMPENS J., TOMASSEN H. B. M. & BERENDSE F. 2003. Expansion of *Sphagnum fallax* in bogs: striking the balance between N and P availability. *J. Bryol.* 25: 83-90.
- MATUŁA J., WOJTUŃ B., TOMASZEWSKA K. & ŻOŁNIERZ L. 1998. Charakterystyka ekologiczna niektórych zespołów roślinnych torfowisk Sudetów. *Geoekologiczne problemy Karkonoszy* 1: 277-287.
- MATUSZKIEWICZ W. 2001. Przewodnik do oznaczania zbiorowisk roślinnych Polski. In: J. B. FALIŃSKI (ed.). *Vademecum Geobotanicum* 3, 537 pp. Wyd. Nauk. PWN, Warszawa.
- MIREK Z., PIĘKOŚ-MIRKOWA H., ZAJĄC A. & ZAJĄC M. 2002. Flowering plants and pteridophytes of Poland. A checklist. In: Z. MIREK (ed.). *Biodiversity of Poland* 1, 442 pp. W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków.
- MISERERE L., MONTACCHINI F. & BUFFA G. 2003. Ecology of some mire and bog plant communities in the Western Italian Alps. *J. Limnol.* 62(1): 88-96.
- NAVRÁTILOVÁ J. & NAVRÁTIL J. 2005. Vegetation gradients in fishpond mires in relation to seasonal fluctuations in environmental factors. *Preslia* 77: 405-418.
- OCHYRA R., ŻARNOWIEC J. & BEDNAREK-OCHYRA H. 2003. Census Catalogue of Polish Mosses. In: Z. MIREK (ed.). *Biodiversity of Poland* 3, 372 pp. Polish Academy of Sciences, Institute of Botany, Kraków.
- PEŁECHATY M. 2004. Can reed stands be good indicators of environmental conditions of the lake littoral? A synecological investigation of *Phragmites australis* – dominated phytocoenoses. *Pol. J. Environ. Studies* 13(2): 177-183.
- PIETSCH W. 1976. Vegetationsentwicklung und wasserchemische Faktoren in Moorgewässern verschiedener Naturschutzgebiete der DDR. *Arch. Naturschutz u. Landschaftsforsch. Berlin* 16(1): 1-43.
- SIEPAK J. (ed.). 1992. Fizyczno-chemiczna analiza wód i gruntów. 193 pp. UAM, Poznań.
- SUCCOW M. 1974. Vorschlag einer systematischen Neugliederung der mineralbodenwasserbeeinflussten wachsenden Moorvegetation Mitteleuropas unter Ausklammerung des Gebirgsraums. *Feddes. Repert.* 85: 57-113.
- SZMEJA J. 2000. Tendencies of changes in flora and vegetation structure of Pomeranian lakes under the influence on humic substances. In: B. JACKOWIAK & W. ŻUKOWSKI

- (eds.). Mechanisms of Anthropogenic Changes of the Plant Cover. Publications of the Department of Plant Taxonomy of the Adam Mickiewicz University in Poznań 10: 85-98. Bogucki Wyd. Nauk., Poznań.
- TAHVANAINEN T. & TUOMALA T. 2003. The reliability of mire water pH measurements – a standard sampling protocol and implications to ecological theory. *Wetlands* 23: 701-708.
- TIMMERMANN T. 1993. Die Meelake – Vegetation und Genese eines Verlandungsmoores in Nordostbrandenburg. *Verh. Bot. Ver. Prov. Brandenburg* 126: 25-62.
- WILLIAMSON C. E., MORRIS D. P., PACE M. L. & OLSON O. G. 1999. Dissolved organic carbon and nutrients as regulators of lake ecosystems: resurrection of more integrated paradigm. *Limnol. Oceanogr.* 44(3): 795-803.
- WOJCIECHOWSKI I. 1970. Mechanizm dystrofizacji jezior sosnowickich. In: VIII Zjazd Hydrobiologów Polskich w Białymstoku, 16-20 września 1970. Streszczenia referatów, p. 180. Polskie Towarzystwo Hydrobiologiczne, Warszawa.
- WOJCIECHOWSKI I. 1990. The trends of lake evolution due to allochthonous humic substances. In: L. BURCHARDT L. (ed.), IX Symposium Phycological Section Polish Botanical Association, International Symposium "Evolution of freshwater lakes", Poznań, 16-22 May 1990, Poland. Uniwersytet im. A. Mickiewicza w Poznaniu, seria Biologia 43: 87-89.
- WOJCIECHOWSKI I. & GÓRNIAK A. 1990. Influence of the brown humic and fulvic acids originating from nearby peat bogs on phytoplankton activity in the littoral of two lakes in Mid-Eastern Poland. *Verh. Internat. Verein. Limnol.* 24: 295-297.
- Woś A. 1994. *Klimat Niziny Wielkopolskiej*. 192 pp. Wyd. Nauk. UAM, Poznań.
- WOJTUŃ B. 2006. Peat mosses (*Sphagnaceae*) in mires of the Sudetes Mountains (SW Poland): a floristic and ecological study. University of Agriculture, Wrocław, Poland, 225 pp.