

Ecosystems with the endemic argan tree *Sideroxylon spinosum* L.: their flora and fauna (Algerian Sahara)

Réda Kechairi^{1*}, Zohra Benaouf², Mohammed Ould Safi³, Ahmed Megharbi⁴
& Benamar Benmahiou⁵

¹Department of Ecology and Environment, Laboratory of Ecology and Management of Natural Ecosystems, Faculty of Natural and Life Sciences, Earth and Universal Sciences, Abou Bekr Belkaïd University, Pole 2 of the ring road, Mansoura, BP 13000, Tlemcen, Algeria; ORCID: RK <https://orcid.org/0000-0002-3477-1132>

²Faculty of Biology, University of Mustapha Stambouli Mascara, Algeria; ORCID: ZB <https://orcid.org/0000-0002-5253-435X>

³INRF research station – Adrar, National Institute of Forestry Research, Algeria; ORCID: MOS <https://orcid.org/0000-0002-6717-5302>

⁴Biological Sciences Department, Laboratory of Environment and Sustainable Development, Relizane University, Algeria; ORCID: AM <https://orcid.org/0000-0002-7056-2207>

⁵Department of Forest Resources, Laboratory of Conservatory Management of Water, Soil and Forests and Sustainable Development, Faculty of Natural and Life Sciences, Earth and Universal Sciences, Abou Bekr Belkaïd University, Pole 2 of the ring road, Mansoura, BP 13000, Tlemcen, Algeria; ORCID: BB <https://orcid.org/0000-0002-5620-3742>

* corresponding author (e-mail: kechairir79@gmail.com)

Abstract. The Tindouf argan groves in the south-western Algerian Sahara, beside *Sideroxylon spinosum* L. [syn. *Argania spinosa* (L.) Skeels] have a rich flora and fauna. This study was aimed to assess their species composition. During floristic surveys we recorded 79 plant species, 67 genera, and 31 families. Among them, the Brassicaceae and Fabaceae are the most represented families (12.7% and 11.4% of the flora, respectively), followed by the Asteraceae and Zygophyllaceae (8.9% each). The flora includes 18 rare species (23%) and 16 endemic taxa (20.3%). The Saharo-Sindian element dominates, accounting for 24.1%, compared with 11.4% for the Saharo-Mediterranean and the Mediterranean elements each. The argan tree grows in steppe-like plant formations, including pastures, of which 67% are of the desert type. The flora is composed mostly of phanerophytes (19.3%), nanophanerophytes (11.6%), and chamaephytes (35.9%). The argan tree is present in various small facies, along wadi beds, on sandy, rocky, and gravelly substrates. We recorded 25 species of birds, 17 mammals, 9 reptiles, 1 amphibian, and 15 arthropods, mostly insects. Consequently, the remarkable biological richness found within the argan ecosystems of Tindouf in the Algerian Saharan context deserves the attention of international bodies concerned with nature conservation in order to protect it as a biosphere reserve.

Key words: *Sideroxylon spinosum*, argan grove, biodiversity, flora, fauna, Sahara, Algeria

1. Introduction

The argan tree *Sideroxylon spinosum* L. [syn. *Argania spinosa* (L.) Skeels] is a major tree component of pre-steppe forests, with xeric determinism on the southern margins of the Mediterranean region (Dupont 1993; Quézel 1999). Unfortunately, the species is currently threatened with extinction (Chouaki *et al.* 2006). Abundant argan trees grow naturally in some arid and semi-arid zones, where it plays an irreplaceable role in the ecological balance and in the preservation of biodiversity (Charrouf 1998). It is the northernmost repre-

sentative of a family that includes hardly any tropical species (Seigue 1985). In many places, it is reduced to mediocre, overgrazed bushes (Msanda *et al.* 2005). The fragility of the ecosystems with *S. spinosum* makes them very vulnerable (Errouati 2005) to 2 major risks: (1) the threat from climate-related disturbances that affect their biotope, which make their survival increasingly difficult; and (2) the increasing human impact, often for socio-economic reasons (Seigue 1985).

From the phytosociological point of view, the argan tree forms several plant associations whose ecological and especially floristic and physiognomic features give

the landscape exceptional originality (Barbero *et al.* 1982; Peltier 1982; Benabid & Fennane 1994; Benabid 2000; Aka Koutoua 2006). The landscape is considerably more complex (Boudy 1950; Feddi *et al.* 2011), containing plant species characteristic of the wooded savannah (Benabid 2000), where the argan tree is part of a pseudo-steppe formation (Zahran 2010).

The argan tree is well adapted to drought conditions in the Algerian western Sahara. The argan groves of Tindouf have a structure of coppice on stumps, with an average height of 4.7 m (Kechairi 2021a). On rocky terrain, the argan plant formation is found in sandy, rocky, and gravelly wadi beds. It covers 56 000 hectares, between 320 and 630 m in altitude (Kechairi & Abdoun 2016). The area of argan groves is characterized by extreme climatic conditions and a dry period that extends all the year round. Under the severe aridity conditions the argan tree adapts well to the hydrography of Hamada of Tindouf in the presence of underflows, where additional water supply is accumulated below the substrate of dry wadi beds (Kechairi 2021b).

The aim of this work in the western Algerian Sahara was to study the previously poorly understood biological diversity within our study area, through floristic and faunistic surveys. Especially the fauna of the argan groves of Tindouf, was mostly not inventoried earlier (Lakhdari & Kechairi 2009).

2. Material and methods

2.1. Study area

Field research on the biodiversity of argan groves in Tindouf Province in the extreme west of the Algerian

Western Sahara was carried out in the north-west of the plateau of Hamada of the river Draa. Taking into account the scattered distribution of groves in the north-west, the area is limited in the north and west by the Moroccan borders, in the south by the Hamada of Tindouf and in the east by the Khebi Wadi (Fig. 1). Our study area is located between 28°17'38.7384"N, 8°40'3.6552"W and 28°45'18.7524"N, 8°2'24.2988"W. The argan groves are situated far from the urban community, about 90 km north-west of the town of Tindouf. It is found between 254 and 634 m above sea level, occupying an area of 50 670 hectares (Kechairi & Abdoun 2016; Kechairi 2018).

Climatic conditions in the study area are generally severe. The maximum temperature reaches 50°C and the dry period extends throughout the year (Kechairi 2009). With annual average precipitation of around 33.5 mm (over a period of 34 years), the argan forest is developing in a particularly precarious climatic context (Kaabèche *et al.* 2010). The hydrographic formation on the desert plateau of the Hamada of the Draa has a mainly alluvial supply, and vegetation generally settles in the beds of wadis.

2.2. Methods

In the spring of 2018, during a series of field trips, we investigated 7 representative plots: 5 at the Touaref Bou-aam station, 1 at Merkala, and 1 at Targant. The phytocological analysis of plant associations with argan trees was carried out through a systematic inventory of quadrats of equal length, forming a regular network (Long 1974). Table 1 presents the survey model adopted in the quadrats. Sampling within the quadrats




Fig. 1. Location of the Tindouf argan groves

Table 1. Floristic survey model adopted with R quadrats (meshes) for each plot

Species	R_1	R_2	R_3	R_4	R_{10}
Species 1	1	0	0	1	-	-	0
Species 2	0	0	1	1	-	-	1
Species (n)	0	0	1	1	-	-	0

was recorded as species presence/absence, according to Quézel (1965) and Braun-Blanquet (1964). Each plot included 10 quadrats (in total, 70 quadrats) and each quadrat was 100 m long, so the length of each plot was 1 km. However, quadrat width was variable, depending on the width of the wadi beds (Fig. 2).

The herbarium specimens and the photos taken were used for the identification of species collected at the

Ecology and Environment Laboratory of the University of Science and Technology Houari Boumediene in Algeria. The identification, classification, rarity assessment, and phytoecological analysis of the species inventoried were based on the works of Quézel and Santa (1962, 1963), Ozenda (1977), Martínez *et al.* (2010), and current updates [<https://science.mnhn.fr/institution/mnhn/search>]. Floristic diversity was expressed in accordance with Daget and Poissonet (1991).

The faunistic list of argan groves was compiled after having identified what is well observed, based on photos, specimens, niches, footprints and excrements, bird feathers, etc. The inventoried fauna was identified according to Cuzin (1998), Ahmim (2019), and the faunistic identification sheets at the Forest Conservation Department in Tindouf.

3. Results

3.1. Flora and vegetation

Our study shows that the Tindouf argan groves are home to 79 plant species (Appendix 1). Downstream the acacias dominate the facies. *Acacia tortilis* subsp. *radiata* and *Faidherbia albida* (syn. *Acacia albida*) are mostly located at the mouth of Touaref Bou-aam, followed by stratified facies, which are found on mounds where some species competing with the argan tree are found, notably *Rhus tripartita* and *Ziziphus lotus* (Fig. 3a).

At mid-slope, on the rocky sand of the wadi beds, the main argan tree groups are found in facies with *Zilla macroptera*, *Farsetia aegyptiaca*, *Echium pycnanthum*, and *Foleyola billoti*. On sandy beds, the main competitor affecting the argan tree remains *Cocculus pendulus* (Fig. 3, b1); the plant communities are dominated by *Psoralea plicata* and *Retama retam* (Fig. 3, b2). There are also patches with *Pergularia tomentosa*, *Echium panicum*, and *Launaea arborescens*. In the rocky upstream quadrates, the vegetation is made up of argan tree groups with *Gymnocarpus decander* and *Farsetia aegyptiaca*, while in the lowland oasis of Targant, with *Tamarix* sp., *Nerium oleander*, *Atriplex halimus*, *Juncus maritimus*, etc. Interestingly, *Faidherbia albida* in the Tindouf argan grove at the mouth of Touaref Bou-aam reaches a remarkable height of over 25 m (Fig. 3c).




Fig. 2. Shapes of sampling quadrats in the wadi beds
 Explanations: a – minor wadi bed, b – major wadi bed




Fig. 3. Plant associations with argan trees on sandy beds

Explanations: a – facies with *Faidherbia albida*, *Ziziphus lotus*, and *Rhus tripartita*, b1 – *Cocculus pendulus*, b2 – *Retama retum*, c – tall *Faidherbia albida* trees at the mouth of Touaref Bou-aam

3.1.1. Correspondence analysis

The analysis was carried out on the basis of the input matrix formed by 3 columns corresponding to the 3 stations (groves) studied and 79 rows representing the species encountered (Appendix 1). The analysis of the floristic surveys of 3 stations shows their grouping into

6 sets, of which 3 are the main ones, relating to each station (Gr1, Gr2, and Gr3). Two intermediate sets are made up of 15 species shared by Merkala and Touaref Bou-aam (Gr4) and 2 species shared by Touaref Bou-aam and Targant (Gr5). The set shared by all the stations (Gr6) consists of 8 species (Fig. 4). The factorial plane (F1x F2) forms 86.6% of the total inertia, with a




Fig. 4. Analysis of the floristic data by projection of the surveys on the factorial plane (F1x F2) of correspondence analysis

Table 2. Eigenvalues of the correlation matrix (correspondence analysis) of floristic data from the Tindouf argan groves

Own value	1.9001	0.6983	0.4015
Proportion	0.633	0.233	0.134
Cumulative	0.633	0.866	1.000

Table 3. Specific contributions of the correlation matrix (correspondence analysis) of floristic data from the Tindouf argan groves


Variable	PC1	PC2
Touaref Bou-aam	0.616	-0.314
Merkala	0.502	0.863
Targant	-0.607	0.395

contribution of 63.3% for axis 1 and 23.3% for axis 2 (Tables 2-3). The point cloud consists of the 6 sets of species. Gr1, marginalized on the negative side of axis 1, is characterized by the floristic composition present in the Targant station, dominated by halophytes due to the halophilic nature of the substrate and the presence of gueltas and springs. The main plant families in this station are the Zygophyllaceae, Salsolaceae, Brassicaceae,

and Tamaricaceae, and the main genera: *Salsola* and *Atriplex*. On the negative side of axis 2, the species of Gr2, characteristic of the Touaref Bou-aam station, are generally dominated by the Fabaceae. On the positive side of axis 2, in Gr3, the therophyte species specific to the Merkala station stand out. Gr4, intermediate between Merkala station and Touaref Bou-aam, is formed by alluvial species, dominated by chamaephytes. Gr5, intermediate between the Targant station and Touaref Bou-aam, is formed by 2 species. Gr6 includes the species that are shared by the 3 stations, namely the key species *Sideroxylon spinosum* and *Acacia tortilis* subsp. *raddiana*.

3.1.2. Plant families

We assessed contributions of 31 botanical families to the flora of the study area (Fig. 5). The 2 most represented families – Brassicaceae and Fabaceae. The families Asteraceae and Zygophyllaceae contribute 8.9% each, followed by the family Amaranthaceae, Poaceae, Apocynaceae, Solanaceae and Boraginaceae (3.8% each), Tamaricaceae, Polygonaceae, and Caryophyllaceae (2.5% each). The remaining 19 families are represented by single species (1.3% each), which shows the high diversity of botanical taxa in the Tindouf argan groves.

**Fig. 5.** Contributions of botanical families to the flora of the argan groves

3.1.3. Degrees of rarity

Out of the 79 plant species recorded in the Tindouf argan groves, 3 taxa are fairly rare (AR) and 11 taxa are rare (R), and 4 taxa are very rare and thus threatened with extinction (RR). In contrast, 37% of species are common (C), 15% very common (CC) and 24% fairly common (AC) (Fig. 6).

3.1.4. Biogeographical elements

In the Tindouf argan groves, endemism is represented by 16 plant taxa (20.3%). The biogeographical distribution of the other species is dominated by the Saharan-Sindian element, which includes 19 taxa (24.1%), characteristic of the Saharan rim (Capot-Rey 1953; Quézel 1965; Barry & Celles 1973). The Saharo-




Fig. 6. Rarity spectrum of the floristic list
Explanations: C – common, AC – fairly common, CC – very common, R – rare, RR – very rare, AR – fairly rare, CCC – extremely common




Fig. 7. Contributions of biogeographic groups to the total number of plant species identified in the argan groves of Tindouf

Mediterranean and Mediterranean elements constitute 11.4% each. The cosmopolitan element accounts for 6.3%, while the Saharo-tropical and the Saharo-Sindian-Mediterranean for 5.1% and 3.8%, respectively. Contributions of other elements to the flora of Tindouf argan groves are low (Fig. 7).

3.1.5. Plant life-forms

Fig. 8 shows the predominance of chamaephytes in Merkala (14 taxa, i.e. 40.4%) and Touaref Bou-aam (19 taxa, i.e. 43.8%). In the Targant station they constituted 23.7% due to the nature of the lowland. The distribution of plant life-forms in the flora of our study region is completely natural. Their average contributions at the 3

sites form the following sequence: chamaephytes (Ch) > therophytes (Th) > phanerophytes (Ph) > nanophanerophytes (NP) > geophytes (Ge) > hemicryptophytes (He).

3.2. Types of biotopes

Our argan groves are mainly represented on the rocky plateau reg on 3 wadi bed substrates: sandy, gravelly, and rocky (Fig. 9). The wadi beds are often loaded with alluvial and wind-blown inflow soils. However, the geomorphological nature of the Hamada is characterized by a rocky biotope, which represents 39.1% in Touaref Bou-aam and 30.6% in Merkala. The colluvial deposits which are frequent in Targant (33.3%), are linked to the rugged nature of the cliffs in the resort. The sandy




Fig. 8. Contributions of plant life-forms to the total number of plant species at the 3 study stations

Explanations: Th – therophyte, G – geophyte, H – hemicryptophyte, Ch – chamaephyte, NPh – nanophanerophyte, MPh – megaphanerophyte




Fig. 9. Plant succession on (a) pebble reg substrate; (b) sandy wadi bed




Fig. 10. Contributions of various biotopes to the total area of the 3 argan groves

substrate is more common in Touaref Bou-aam (33.3%) than in the other stations (Fig. 10).

3.3. Types of pasture

The distribution of species by geographical types increases the diversity of pasture types, both for herbivorous wild animals and for nomadic herds. Desert grazing represents 67% in our study region (Fig. 11).

3.4. Animal diversity

The argan tree is home to a great diversity of animals and plants, including nearly 300 invertebrate taxa (Naamani 2004). Thus the Tindouf nature reserve not only protects *Sideroxylon spinosum* but also provides a favourable shelter for many vertebrates and invertebrates (Fig. 12). As a result, the fauna as a whole benefits from the protection offered by the tree (Er-Rafia 1975), which




Fig. 11. Spectrum of the various types of pastures in the study region




Fig. 12. Some animal species of the Tindouf argan groves

Explanations: a – *Mellivora capensis leucanota*, b – *Pelophylax saharicus*, c – *Ptyodactylus oudrii*, d – *Equus asinus*, e – *Uromastix acanthinurus*, and mammal footprints or shelters under argan trees: (f1, f2) *Vulpes vulpes* footprints and its shelter, G1 – squirrel footprints, G2 – wild rabbit shelter (original photos by R. Kechairi taken in the Tindouf argan groves)

is also an important habitat for birds (Vernon *et al.* 2005). We recorded 25 species of birds, 17 mammals, 9 reptiles, 1 amphibian, and 15 arthropods, mostly insects (Table 4).

The environmental importance of this landscape is remarkable. Firstly, it is a breeding area for endangered wildlife species. Secondly, the groves accumulate and restore the soils. Thirdly, the area is a transition station for migratory birds through the small marshes north of the argan groves (Tafegoumet and Oum El-Achar). Fourthly, it allows safe passage of dorcas gazelles and Barbary sheep away from the villagers and the haters along the southern slope of the river Draa and the north of Ouarkziz crater.

4. Discussion

This paper presents the first study of the floristic and faunistic richness of argan groves in the west of the Algerian Sahara. The delimitation of the distribution of *Sideroxylon spinosum* and the comprehensive inventory of plant and animal species were launched by Kechairi (2009). In the same circumstances, an ecological description, presentation of the physical environment, and an imprecise delimitation of its distribution area

were carried out by Benkheira (2009). Subsequently, Kaabèche *et al.* (2010) focused on a phytosociological analysis. However, Kechabar *et al.* (2013) detailed the distribution of argan trees in the main wadi beds.

Through subjective sampling we compiled a floristic list of 79 species divided into 31 botanical families. Based on the scale of Daget and Poissonet (1991), the flora of the region is very rich. The argan tree is found in various plant associations in the form of small facies along the El-Ma wadi and its tributaries. The distribution of the facies is influenced by the nature of the lithological substrate of the wadi beds.

The basic structure of the argan grove is a coppice forest, with a dominance of dense and thorny bushes (Kechairi 2021a), in a desert savannah-type plant formation with acacias and *Panicum turgidum*, characteristic of the Sahara in North-West Africa (Quézel & Barbero 1993) and used for grazing (Boughani *et al.* 2009). On alluvial beds, we have essentially the same formation observed in south-western Morocco: *Periploco laevigatae-Agranietum spinosae*, which grows on limestone and dolomite (Msanda 1993; Peltier & Msanda 1995).

The argan ecosystem is distinguished from the surrounding desert environment by the presence of vigorous

Table 4. List of animal species recorded in the Tindouf argan groves

Animal group	Species
Birds	<i>Accipiter nisus</i> (Linnaeus, 1758) Eurasian Sparrowhawk; <i>Alaemon alaudipes</i> (Desfontaines, 1789) Greater Hoopoe-lark; <i>Ammomanes deserti</i> (Lichtenstein, 1823) Desert Lark; <i>Ammomanes cinctura</i> (Gould, 1839) Bar-tailed Lark; <i>Buteo rufinus</i> (Cretzschmar, 1827) Long-legged Buzzard; <i>Calandrella brachydactyla</i> (Leisler, 1814) Greater Short-toed Lark; <i>Caprimulgus aegyptius</i> (Lichtenstein, 1823) Egyptian Nightjar; <i>Chlamydotis undulate</i> (Jacquin, 1784) African houbara; <i>Ciconia ciconia</i> (Linnaeus, 1758) White stork; <i>Corvus ruficollis</i> (Lesson, 1831) Brown-necked Raven; <i>Cursorius cursor</i> (Latham, 1787) Cream-coloured Courser; <i>Galerida cristata</i> (Linnaeus, 1758) Crested Lark; <i>Gyps fulvus</i> (Hablizl, 1783) Griffon Vulture; <i>Hirundo rustica</i> (Linnaeus, 1758) Barn Swallow; <i>Leiothlypis peregrina</i> (Wilson, 1811) Tennessee Warbler; <i>Motacilla alba</i> (Linnaeus, 1758) White Wagtail; <i>Neophron percnopterus</i> (Linnaeus, 1758) Egyptian vulture; <i>Oenanthe leucopyga</i> (Brehm, 1855) White-crowned; <i>Phoenicurus moussieri</i> (Olphe-Galliard, 1852) Moussier's Redstart; <i>Phylloscopus humei</i> (Brooks, 1878) Hume's Leaf-warbler; <i>Pterocles alchata</i> (Linnaeus, 1766) Pin-tailed Sandgrouse; <i>Scotocerca inquieta</i> (Cretzschmar, 1827) Streaked Scrub-warbler; <i>Streptopelia decaocto</i> (Fridvaldszky, 1838) Eurasian Collared-dove; <i>Turdoides fulvus</i> (Desfontaines, 1789) <i>Fulvous Babbler</i> ; <i>Upupa epops</i> (Linnaeus, 1758) Common Hoopoe
Mammals	<i>Ammotragus lervia</i> subsp. <i>sahariensis</i> (Rothschild, 1913) Barbary sheep (observed on Ouarkiz crater and in Targant); <i>Atelerix algirus</i> (Lereboullet, 1842) Algerian hedgehog; <i>Atlantoxerus getulus</i> (Linnaeus, 1758) Barbary ground squirrel (major pest of argan seeds); <i>Camelus dromedarius</i> (Linnaeus, 1758) Arabian camel; <i>Canis anthus</i> (F.Cuvier, 1820) African Golden Wolf; <i>Equus asinus</i> (Linnaeus, 1758) Wild Donkey; <i>Felis sylvestris</i> (Schreber, 1777) Wild Cat; <i>Gazella dorcas</i> (Linnaeus, 1758) Dorcas Gazelle; <i>Gerbillus campestris</i> (Loche, 1867) North African gerbil; <i>Hystrix cristata</i> (Linnaeus, 1758) Crested porcupine; <i>Jaculus jaculus</i> (Linnaeus, 1758) Lesser jerboa; <i>Lepus capensis</i> (Linnaeus, 1758) Cape hare; <i>Mellivora capensis</i> (Schreber, 1776) Honey badger; <i>Meriones crassus</i> (Sundevall, 1842) Sundevall's jird; <i>Psammomys obesus</i> (Cretzschmar, 1828) Fat sand rat; <i>Vulpes vulpes</i> (Linnaeus, 1758) Red fox; <i>Vulpes zerda</i> (Zimmermann, 1780) Fennec fox
Reptiles	<i>Acanthodactylus boskianus</i> (Daudin, 1802) Bosc's Acanthodactyl; <i>Agama impalearis</i> (Boettger, 1874) Bibron's agama; <i>Cerastes cerastes</i> (Linnaeus, 1758) Desert Horned Viper; <i>Macrovipera mauritanica</i> (Duméril & Bibron, 1848) Sahara Rock Viper; <i>Mesalina olivieri</i> (Audouin, 1829) Olivier's Sand Lizard; <i>Ptyodactylus oudrii</i> (Lataste, 1880) Algerian Fan-fingered Gecko (observed in small water bodies in Targant valley); <i>Trapelus mutabilis</i> (Merrem, 1820) Desert agama; <i>Uromastix acanthinura</i> (Bell, 1825) Bell's Dabb Lizard; <i>Varanus griseus</i> (Daudin, 1803) Desert monitor
Amphibians	<i>Pelophylax saharicus</i> (Boulenger, 1913) syn. <i>Rana saharica</i> (Sahara frog)
Arthropods (arachnids and insects)	<i>Androctonus liouvillei</i> (Pallary, 1924) Black Scorpion; <i>Anthophila</i> sp. (Bee); <i>Buthus atlantis</i> (Pocock, 1889) Atlas Scorpion; <i>Ceratitis capitata</i> (Wiedemann, 1824) Mediterranean fruit fly; <i>Dociostaurus maroccanus</i> (Thunberg, 1815) Moroccan locust; <i>Galeodes olivieri</i> (Simon, 1879) Sun spider; <i>Harmonia</i> sp. (Ladybird); <i>Hottentotta gentili</i> (Pallary, 1924) black Scorpion; <i>Heteronychus arator</i> (Fabricius, 1775) African Black Beetle; <i>Odonata</i> (Fabricius, 1793) Dragonfly; <i>Messor</i> sp. (Forel, 1890) Harvester ant; <i>Reticulitermes</i> sp. (Holmgren, 1913) Termite; <i>Schistocerca gregaria</i> (Forskål, 1775) Desert locust; <i>Trithemis pallidinervis</i> (Kirby, 1889) Dragonfly; <i>Vanessa cardui</i> (Linnaeus, 1758) Butterfly; <i>Vespa</i> (Thomson, 1869) Ground Yellowjackets

argan trees over 10 m high and *Faidherbia albida* over 25 m high in Wadi El-Ma (Kechairi 2021a). This confirms what Ozenda said (1991), that trees are far from being absent from the Saharan flora and that there are at least 20 arborescent species in the north and centre of the Sahara. Most of them can reach a size comparable to those of trees of temperate zones (Quézel & Barbero 1993).

The most continental in the whole biogeographic classification of North-West Africa, the floristic elements in our study area are made up of similar types already cited by Peltier (1982): tropical, Saharan, and Mediterranean, but the Macaronesian element is absent. The Mediterranean element confirms that the region has a great floristic originality (Chevalier 1943).

The majority of Saharan trees are of tropical origin (Quézel & Barbero 1993), but in our study *Sideroxylon spinosum*, *Acacia* spp., and *Ziziphus lotus* dominate in the canopy. Where the argan tree is present, endemism is high (Zahran 2010) and indeed endemic species constitute 20.3% of the flora of the studied plant formation.

Considering plant life-forms, chamaephytes dominate with a frequency of around 40% in Touaref Bou-aam and Merkala. Unfortunately, these sites are currently considered an open field for the herds of Sahrawi nomads. Therophytes, which are a sign of aridity of the environment (Négre 1966), have a frequency of 34.2% in the border region of Targant, not accessible to herds (Kechairi 2021a).

It is alarming that the increase in anthropogenic and environmental disturbances, including climate change (Moukrim *et al.* 2019), negatively affect argan forest ecosystems. Therefore, long-term efforts must be taken within the framework of sustainable development to protect these ecosystems, threatened with extinction if they do not receive enough attention, not only at the national level, but also at the global level.

5. Conclusions

The possible disappearance of the argan groves could lead to a great decrease in biodiversity, including

a reduction in genetic heritage. The ecosystems with argan trees play an irreplaceable role in the ecological balance, with many living organisms (fauna, flora and microflora) directly linked to its presence. Its natural environment at the Hamada of the Draa includes valuable vegetation, occupying the ecological niches on the wadi beds, with specific biological diversity that must be preserved in order to maintain the authentic habitat for threatened animal species, such as the fennec fox, spiny-tailed lizard, and houbara bustard. Unfortunately the ostrich has already completely disappeared.

Our floristic analysis has shown that the Brassicaceae and Fabaceae are the most represented families (12.7% and 11.4% of the flora, respectively), followed by and the Asteraceae and Zygophyllaceae (8.9% each), while 18 species (23%) are rare and 16 are endemic (20.3%). The Saharo-Sindian element is the major biogeographical group (24.1% of the flora), compared with 11.4% for the Saharo-Mediterranean and the Mediterranean elements each.

In the Algerian Sahara, the Tindouf argan groves are valuable ecosystems, which serve as shelters for wildlife. Indeed, the argan ecosystem is characterized

by a rich flora, represented by 31 families, 67 genera, and 79 species, and we recorded 25 species of birds, 17 mammals, 9 reptiles, 1 amphibian, and 15 arthropods, mostly insects. The argan groves need better protection and conservation. Their degradation could have a considerable impact on biodiversity, inducing its reduction and a drift of the genetic heritage for the tree as well as for other animal, plant or microbial species.

Our argan heritage, which constitutes a valuable ecosystem within the Sahara desert, requires long-term conservation management within the framework of sustainable development. Firstly, by local authorities and services concerned with nature protection at the national level, but also by seeking the support of NGOs through international cooperation.

Author Contributions:

Research concept and design: R. Kechairi

Collection and/or assembly of data: R. Kechairi

Data analysis and interpretation: R. Kechairi, A. Megharbi

Writing the article: R. Kechairi

Critical revision of the article: Z. Benaouf, M. Ould Safi, B. Benmahioul

Final approval of article: R. Kechairi

References

- AHMIM M. 2019. The Wild Mammals of Algeria: Distribution and Conservation Biology. Les Editions du Net, 2019, 978-2312068961.fhhal-02375326f.
- AKA KOUTOUA M. 2006. Ecological, Phytosociological Analysis and evaluation of the balance sheets of Argan tree (*Argania spinosa*) plantations for the regeneration and rehabilitation of its natural ecosystems (Agadir, Taroudant and Tiznit regions). In: DREF/SO. ENFI Salé, Morocco, 113 p.
- BARBERO M., BENABID A., QUÉZEL P., RIVAS-MARTINEZ S. & SANTOS A. 1982. Contribution to the study of the *Acacio-Arganietalia* of south-western Morocco. Doc. Phytosociologique 6: 311-338.
- BARRY J. P. & CELLES J. C. 1973. The problem of bioclimatic and floristic divisions in the Algerian Sahara. *Naturalia monspeliensia*, (1972-1973), Serie botanique 23-24: 5-48.
- BENABID A. 2000. Flore et écosystèmes du Maroc: évaluation et préservation de la biodiversité. 359 pp. Ibis Press. Paris.
- BENABID A. & FENNANE M. 1994. Knowledge of vegetation in Morocco: phytogeography, phytosociology and vegetation series. *Lazaroa* 14: 21-97.
- BENKHEIRA A. 2009. L'Arganeraie algérienne. Bulletin d'information du Projet ALG/00/G35, Conservation de la biodiversité et gestion durable des ressources naturelles, Numéro spécial, pp. 1-15.
- BOUDY P. 1950. Monograph and treatment of the Argan tree. In Monographie et traitements des essences forestières. Larose edn, Paris, France.
- BOUGHANI A., SADKI N., MÉDAILLE F., NEDJRAOUI D. & SALAMANI M. 2009. Floristic and phytogeographic analysis of a region of the Saharan Constantine Atlas, the Ghouffi gorges (Algeria). *Acta Botanica Gallica* 156(3): 399-414. <https://doi.org/10.1080/12538078.2009.10516166>
- BRAUN-BLANQUET J. 1964. Sociology of Plants: Basic Principles of Vegetation Science. Springer, 3th 865 pp. Ed. Vienna-New York.
- CAPOT-REY R. 1953. Les limites du Sahara français. *Trav. Inst. Rech. Sahariennes* 8: 23-48. <https://doi.org/10.3406/ahess.1953.2140>
- CHARROUF Z. 1998. Valorization of Argan tree products for a sustainable management of the arid zones of south-west Morocco. Mohammed V. University. Morocco.
- CHEVALIER A. 1943. Les Sapotacées à graines oléagineuses et leur avenir en culture. *Journal d'agriculture traditionnelle et de botanique appliquée* 23: 97-159. <https://doi.org/10.3406/jatba.1943.1753>
- CHOUAKI S., BESSÉDIK F., CHEBOUTI A., MAAMRI F., OUMATA S., KHELDOUN S., HAMANA M., DOUZENE M., BELLAH F. & KHELDOUN A. 2006. Second national report on the state of plant genetic resources. INRAA/FAO/June.
- CUZIN F. 1998. Propositions pour le plan de gestion du Parc National du Bas Drâa. 73 pp. GTZ Ed.
- DAGET P. & POISSONET J. 1991. Meadows and pastures: Methods of study. Institute of Botany, Montpellier. French, pp. 285-324.
- DUPONT L. M. 1993. Vegetation zones in NW Africa during the brunhes chron reconstructed from marine palynological

- data. *Quaternary Science Reviews* 12(3): 189-202. [https://doi.org/10.1016/0277-3791\(93\)90053-O](https://doi.org/10.1016/0277-3791(93)90053-O)
- ER-RAFIA M. 1975. The Argan grove and its problems. Personal theme ENGREF. Nancy, 38 p.
- ERROUATI A. 2005. Problem of assisted regeneration and reforestation based on *Argania spinosa* in the region of the Amsitten forest massif (Province of Essaouira). Phd thesis, ENFI, Salé, Morocco, 142 p.
- FEDDI N., FAUQUETTE S. & SUC J.-P. 2011. Plio-Pleistocene history of plant ecosystems in the south-western Mediterranean: contribution of pollen analysis from two surveys in the Alboran Sea. *Geobios* 44(1): 57-69. <https://doi.org/10.1016/j.geobios.2010.03.007>
- KAABÈCHE M., BENKHEIRA A. & DE FOUCAULT B. 2010. L'arganeraie d'Algérie: structure, écologie, syntaxonomie, dynamique. *Acta Botanica Gallica* 157(3): 563-572. <https://doi.org/10.1080/12538078.2010.10516231>
- KECHAIRI R. & ABDOUN F. 2016. Cartographic status of the Argan tree *Argania spinosa* (L.) Skeels (Sapotaceae) in North-West Africa (Algeria and Western Sahara). *International Journal of Environmental Studies* 73(2): 286-293. <https://doi.org/10.1080/00207233.2016.1148448>
- KECHAIRI R. 2009. Contribution to the ecological study of the Argan tree *Argania spinosa* (L.) skeels in the region of Tindouf (Algeria). F.S.B. University USTHB Algiers, 75 p.
- KECHAIRI R. 2018. Étude de l'Arganeraie de Tindouf : État des lieux, contraintes et perspectives de son développement. Département des ressources forestières. Faculté SNV/STU. Université de Tlemcen, Algérie, 196 p.
- KECHAIRI R. 2021a. Structure élémentaire de l'arganeraie *Argania spinosa* L. Skeels de Tindouf (Sahara occidental algérien) *Ecologia Mediterranea* 27(2): 73-84. <https://doi.org/10.3406/ecmed.2021.2131>
- KECHAIRI R. 2021b. Argan tree hydrography water sources water compensation Tindouf, Algeria. *Algerian Journal of Arid Environment* 11(1): 90-97.
- KECHAIRI R., OULD SAFI M. & BENMAHIOUL B. 2018. Étude comparative de deux plantations d'*Argania spinosa* (L.) Skeels (Sapotaceae) dans le Sahara Occidental Algérien (Tindouf et Adrar). *International Journal of Environmental Studies* 75(2): 294-308. <https://doi.org/10.1080/00207233.2017.1360601>
- KECHEBAR M., KAROUNE S., BELHAMRA M. & RAHMOUNE C. 2013. Étude structurale des peuplements d'arganier (*Argania spinosa*) en Algérie. *Journal Algérien des Régions Arides* 12(1): 46-54.
- LAKHDARI A. & KECHAIRI R. 2009. La faune de l'Arganier de Tindouf (Algérie), In: Actes du Séminaire International sur la Biodiversité Faunistique en Zones Arides et Semi-arides, Université d'Ourglia, Algérie, pp. 207-211.
- LONG G. 1974. Diagnostic phytoécologique et aménagement du territoire. I: Principes généraux et méthodes. Masson édit., Paris. Vol. 1. 252 p.
- MARTÍNEZ I. B., ÁLVAREZ M. E. R., SOTOMAYOR S. P. & MUSTAPHA R. S. 2010. Sahara Occidental. Plantas y Usos. Estudio etnobotánico del Sahara Occidental: Usos y costumbres del pueblo saharaui relacionados con los recursos vegetales. *Monografías Botánica Ibérica* 4: 1-117.
- MOUKRIM S., LAHSSINI S., RHAZI M., ALAOUI H.M., BENABOU A., WAHBY I., EL MADIHI M., ARAHOU M. & RHAZI L. 2019. Climate change impacts on potential distribution of multipurpose agro-forestry species: *Argania spinosa* (L.) Skeels as case study. *Agroforestry systems* 93(4): 1209-1219. <https://doi.org/10.1007/s10457-018-0232-8>
- MSANDA F. 1993. Ecology and cartography of plant groups in Anzi (Anti-Western Atlas, Morocco) and contribution to the study of the genetic diversity of the Argan tree *Argania spinosa* (L.) Skeels. Univ. Joseph Fourier, Grenoble 1. France, 116 p.
- MSANDA F., EL ABOUDI A. & PELTIER J.-P. 2005. Biodiversity and biogeography of the Moroccan Argan forest. *Cahiers Agricultures* 14(4): 357-364.
- NAAMANI K. 2004. Study of the entomofauna of the Argan tree (*Argania spinosa* L. Skeels) and the bio-ecology of its pest *Ceratitis capitata* Wiedemann (Diptera: Tephritidae) in the region of Essaouira (Morocco). Thèse de Doctorat. Département de Biologie. Faculté des Sciences Semlalia, Marrakech, Maroc, 223 p.
- NÉGRE R. 1966. Therapists. *Mem. Soc. Bot. France* pp. 92-108. <https://doi.org/10.1080/00378941.1966.10838477>
- OZENDA P. 1991. Flora of the Sahara. 3rd Edition. CNRS, Paris, 662 p.
- PELTIER J. P. 1982. Vegetation in the catchment area of the Souss wadi (Morocco). Faculty of Science and Medicine of Grenoble, 201 p.
- PELTIER J. P. & MSANDA F. 1995. Diversité et originalité de la steppe à *Euphorbia officinarum* L. subsp. *echinus* (HOOKER fil. et COSSON) VINDT du sud-ouest Marocain. *Feddes Repertorium* 106(3-4): 215-229. <https://doi.org/10.1002/fedr.19951060316>
- QUÉZEL P. 1965. La végétation du Sahara: du Tchad à la Mauritanie. *Geobotanica selecta* 2: 1-333.
- QUÉZEL P. 1999. Large vegetation structures in the Mediterranean region: Determining factors in their post-glacial establishment. *Geobios* 32(1): 19-32.
- QUÉZEL P. & BARBERO M. 1993. Variations climatiques au Sahara et en Afrique sèche depuis le Pliocène: enseignements de la flore et de la végétation actuelles. *Bulletin d'écologie* 24: 191-202.
- QUÉZEL P. & SANTA S. 1962, 1963. Nouvelle flore de l'Algérie et des régions désertiques méridionales. Vol. 1-2. Éditions du CNRS, France.
- SEIGUE A. 1985. The circummediterranean forest and its problems. Maisonneuve et Larose Édit., Paris. France.
- VERNON R., THÉVENOT M., BERGIER P. & ROUSSEAU E. 2005. Argan woodland: an important bird habitat in Morocco. *Bull. ABC* 12(2): 134-146. <https://doi.org/10.5962/p.309751>
- ZAHARAN M. A. 2010. Climate-Vegetation: Afro-Asian Mediterranean and Red Sea Coastal Lands. Springer Science & Business Media 4: 219-248. https://doi.org/10.1007/978-90-481-8595-5_3

Appendix 1. Floristic list of the Tindouf argan groves

Family	Species	Plant life-form	Degree of rarity	Biotope	Biogeo-graphical distribution	Stations		
						TFB	MRK	TRG
Adiantaceae	<i>Adiantum capillus-veneris</i> L.	G	AC	Rocks	Med.	0	0	1
Amaranthaceae	<i>Anabasis aretioides</i> Moq. & Coss. ex Bunge	Ch	C	Desert pastures	End.	1	0	0
	<i>Anabasis articulata</i> (Forssk.) Moq.	Ch	C	Desert pastures	Sah.-Sind.	0	0	1
	<i>Atriplex halimus</i> L.	Ch	C	Rockeries	Cosmop.	0	0	1
	<i>Haloxylon scoparium</i> Pomel	Ch	AC	Desert pastures	Sah.-Med.	1	1	1
	<i>Salsola imbricata</i> Forssk.	Ch	R	Desert pastures	Sah.-Sind.	0	0	1
	<i>Salsola tetragona</i> Delile	Ch	AC	Desert pastures	Sah.-Med.	0	0	1
Amaryllidaceae	<i>Pancratium saharae</i> Coss. ex Batt. & Trab. var. <i>chatinianum</i> Batt.	G	AR	Arid pastures and rockeries	Sah.-trop.	1	1	0
Anacardiaceae	<i>Rhus tripartita</i> (Ucria) Large	MPh	AR	Arid pastures and rockeries	Sah.-Med.	1	1	1
Apiaceae	<i>Deverra battandieri</i> (Mayor) Chrték	Ch	AC	Desert pastures	End.	0	1	0
Apocynaceae	<i>Calotropis procera</i> (Aiton) Dryand.	MPh	C	Wadi beds	Sahelo-Sah.	1	0	0
	<i>Nerium oleander</i> L.	NPh	R	Wadi beds, rockeries	Med.	0	0	1
	<i>Pergularia tomentosa</i> L. var. <i>schmidtiana</i> Batt.	Ch	CC	Desert pastures	Sah.-Sind.	1	1	0
	<i>Periploca laevigata</i> subsp. <i>angustifolia</i> (Labill.) Markgr.	NPh	CC	Arid rockeries	Sah.-Med.	0	1	0
Araliaceae	<i>Hedera helix</i> L.	MPh	CC	Wadi beds	Med.	0	0	1
Arecaceae	<i>Phoenix dactylifera</i> L.	MPh	C	Wadi beds	End.	0	0	1
Asteraceae	<i>Aaronsohnia pubescens</i> (Desf.) K. Bremer & Humphries	Th	CC	Desert pastures	N. Afr.	1	0	0
	<i>Anvillea garcinii</i> subsp. <i>radiata</i> (Coss. & Durieu) Eterb	Ch	CC	Stony pastures	End.	1	1	0
	<i>Asteriscus graveolens</i> (Forssk.) Less.	Ch	CCC	Stony pastures	Sah.-Sind.	1	0	0
	<i>Launaea arborescens</i> (Batt.) Murb.	NPh	C	Desert pastures, rockeries	Ibero.-Maur.-Sah.	1	1	1
	<i>Launaea nudicaulis</i> (L.) Hook.f.	Th	CC	Desert pastures	Med. Sah.-Sind.	1	0	0
	<i>Picris coronopifolia</i> (Desf.) DC. sp. <i>albida</i> (Ball) M.	Th	AC	Wadi beds, rockeries	Sah.-Med.	1	0	0
	<i>Rhanterium suaveolens</i> Desf.	Ch	C	Desert pastures	End.	1	0	0
Boraginaceae	<i>Echium humile</i> Desf.	H	C	Desert pastures	Sah.-Med.	1	1	0
	<i>Heliotropium bacciferum</i> Forssk.	Ch	C	Variables	Sah.-Sind.	0	0	1
	<i>Trichodesma africanum</i> (L.) R.Br.	Th	AC	Desert rockeries	Sah.-Trop.	1	1	0
Brassicaceae	<i>Diplotaxis harra</i> (Forssk.) Boiss.	Th	RR	Wadi beds	Med.-Iran-Tur.	0	0	1
	<i>Diplotaxis pitardiana</i> Mayor	Th	R	Wadi beds	End.	0	0	1
	<i>Eremobium aegyptiacum</i> (Spreng.) Asch. ex Boiss.	Th	C	Sands	Sah. Sind.-trop.	1	0	0
	<i>Eruca vesicaria</i> (L.) Cav.	Th	C	Desert pastures	Med.	0	1	0
	<i>Farsetia aegyptiaca</i> Turra	Ch	C	Desert regs	Sah.-Sind.	1	1	0
	<i>Farsetia hamiltonii</i> Royle	Ch	AC	Desert regs	Sah.-Sind.	0	0	1
	<i>Foleyola billotii</i> Mayor	NPh	R	Wadi beds	End.	1	0	1
	<i>Matthiola maroccana</i> Coss. var. <i>puberula</i> Mayor	Th	R	Desert pastures, rockeries	End.	0	1	0
	<i>Morettia canescens</i> Boiss. var. <i>erecta</i> Mayor	Th	C	Desert pastures	Sah.-Sind.	1	0	0
	<i>Zilla spinosa</i> subsp. <i>macroptera</i> (Coss.) Maire & Weiller	Ch	C	Desert rockeries	End.	1	1	1
Capparaceae	<i>Capparis spinosa</i> L.	Ch	AC	Rockeries	Med.- Sah.-Sind.	0	0	1
Caryophyllaceae	<i>Gymnocarpus decander</i> Forssk	Ch	AC	Desert rockeries	Sah.-Sind.	1	1	0
	<i>Paronychia arabica</i> (L.) DC. subsp. <i>tibetica</i> Quézel	Th	AC	Desert pastures	Sah.	1	0	0

Cistaceae	<i>Helianthemum lippii</i> (L.) Dum. Course.	Ch	C	Desert pastures	Sah.-Med.	1	1	0
Colchicaceae	<i>Androcymbium punctatum</i> (Cav.) Baker. var. <i>saharae</i> M.	G	AC	Rocky pastures, Sands	Sah.-Sind.	1	0	0
Convolvulaceae	<i>Convolvulus trabutianus</i> Schw. and Musch.	Ch	R	Desert rockeries	End.	1	0	0
Cucurbitaceae	<i>Citrullus colocynthis</i> (L.) Schrad.	Th	CC	Sands	Trop. Med.	1	1	0
Euphorbiaceae	<i>Euphorbia calyptrate</i> Cosson and DR. var. <i>involutate</i> Batt.	Th	C	Desert pastures	End.	0	0	1
Fabaceae	<i>Acacia ehrenbergiana</i> Hayne	MPh	RR	Wadi beds	Sah.-Med.	1	0	0
	<i>Acacia seyal</i> Delile	MPh	C	Wadi beds	Afr. Sah.-trop.	1	0	0
	<i>Acacia tortilis</i> (Forssk.) Hayne	MPh	C	Wadi beds	Afr. Sah.-trop.	1	1	1
	<i>Astragalus gomboeformis</i> Pomel	Ch	AC	Desert pastures	End.	1	1	0
	<i>Crotalaria saharae</i> Coss.	Ch	C	Wadi beds	End.	1	1	0
	<i>Cullen plicatum</i> (Delile) CH Stirt.	Ch	AC	Wadi beds	Afr. trop.	1	1	0
	<i>Faidherbia albida</i> (Delile) A.Chev	MPh	R	Wadi beds	Afr. trop.	1	0	0
	<i>Lupinus cosentinii</i> Guss.	Th	R	Wadi beds	Sah.-Med.	0	1	0
	<i>Retama raetam</i> (Forssk.) Webb	NPh	C	Sands	Sah.-Sind.	1	1	1
	<i>Monsonia heliotropioides</i> (Cav.) Boiss.	Th	CC	Rockeries	Sah.-Sind.	0	1	0
Juncaceae	<i>Juncus maritimus</i> Lam.	G	C	Gueltas	Cosmop.	0	0	1
Lamiaceae	<i>Salvia aegyptiaca</i> L.	Ch	C	Desert pastures	Sah.-Sind.	1	0	0
Menispermaceae	<i>Cocculus pendulus</i> (JR Forst. & G. Forst.) Diels	NPh	R	Rockeries	Trop.	1	0	1
Plantaginaceae	<i>Plantago amplexicaulis</i> Cav.	Th	AR	Sands	Med.	1	0	0
Poaceae	<i>Aeluropus littoralis</i> (Gouan) Parl.	H	C	Gueltas	Med.	0	0	1
	<i>Cymbopogon schoenanthus</i> subsp. <i>schoenanthus</i> (L.) Spreng.	Ch	C	Wadi beds	Sah.-trop.	1	1	0
	<i>Panicum turgidum</i> Forssk.	Ch	AC	Saharan pastures	Sah.-trop.	1	1	0
Polygonaceae	<i>Phragmites communis</i> Trin.	G	C	Gueltas	Cosmop.	0	0	1
	<i>Schismus arabicus</i> Nees	Th	C	Wadi beds	Med.	0	0	1
	<i>Emex spinosa</i> (L.) Campd.	Th	RR	Wadi beds	Med.	0	0	1
	<i>Rumex simpliciflorus</i> Murb.	Th	C	Desert rockeries	Sah.-Sind.	0	0	1
	<i>Ziziphus lotus</i> (L.) Lam.	MPh	CC	Desert pastures	Med.	1	1	0
Rhamnaceae	<i>Sideroxylon spinosum</i> L., syn. <i>Argania spinosa</i> (L.) Skeels	MPh	RR	Desert pastures, rockeries	End.	1	1	1
Solanaceae	<i>Lycium afrum</i> L.	MPh	AC	Desert pastures, rockeries	Sah.	1	1	1
	<i>Lycium intricatum</i> Boiss.	NPh	R	Rocks	Ibero-Maur.-Sah.	1	1	0
Tamaricaceae	<i>Solanum nigrum</i> L.	Th	R	Rockeries	Cosmop.	0	0	1
	<i>Tamarix articulata</i> Vahl.	Ph	C	Wadi beds	Sah.-Sind.	0	0	1
	<i>Tamarix gallica</i> L.	Ph	CC	Gueltas	Med. Sah.-Sind.	0	0	1
Zygophyllaceae	<i>Fagonia glutinosa</i> Delile	Th	CC	Desert pastures	Sah.-Sind.	0	0	1
	<i>Fagonia latifolia</i> Delile	Th	AC	Desert rockeries	Sah.-Sind.	0	0	1
	<i>Fagonia longispina</i> Batt.	Th	C	Desert pastures, rockeries	End.	0	0	1
	<i>Seetzenia lanata</i> (Willd.) Bullock	Th	AC	Desert pastures	Sah.-Sind.	1	0	0
	<i>Tetraena gaetula</i> subsp. <i>gaetulum</i> (Emb. & Mayor) Beier & Thulin	Th	AC	Gueltas	End.	0	0	1
	<i>Tribulus alatus</i> Forssk.	Th	AC	Desert pastures	Sah.-Sind.	0	0	1
	<i>Tribulus terrestris</i> L.	Th	CC	Desert pastures	Cosmop.	0	0	1

Explanations: stations, TFB – Touaref Bou-aam, MRK – Merkal, TRG – Targant; plant life-form, Ch – chamaephyte, G – geophyte, H – hemicryptophyte, MPh – megaphanerophyte, NPh – nanophanerophyte, Th – therophyte; degree of rarity, AC – fairly common, AR – fairly rare, C – common, CC – very common, CCC – extremely common, R – rare, RR – very rare